



2018 Medicare Nuclear Medicine  
**Reimbursement Information**



## 2018 Medicare Reimbursement Information Lantheus Medical Imaging

If you have questions regarding reimbursement for Lantheus Medical Imaging products, call Randy VanCoughnett at 978-436-7995 or email [randy.vancoughnett@lantheus.com](mailto:randy.vancoughnett@lantheus.com).

### **CPT® – Current Procedural Terminology**

- American Medical Association's five digit numeric codes used to report medical procedures and services.

### **HCPCS - Healthcare Common Procedure Coding System**

- Level I HCPCS codes American Medical Association's Current Procedural Terminology (CPT).
- Level II HCPCS codes alphanumeric five digit codes primarily to identify contrast agents, radiopharmaceuticals, supplies and devices.

### **Q-codes**

- Temporary codes created by Medicare to identify items not assigned a CPT code. Many drugs, supplies and biologicals are assigned Q codes.

### **NDC codes – National Drug Code**

- A unique numeric code to identify drugs. The first segment of numbers identifies the labeler or manufacturer, the second segment identifies the product, and the third identifies the package.

### **HOPPS – Hospital Outpatient Prospective Payment System**

#### **Contents**

Components of Reimbursement	Page 3
Non-HEU derived Tc-99m	Page 4
Medicare Hospital Outpatients	Page 5
NDC Codes	Page 6

**Three Basic Components of Reimbursement: Coding, Coverage and Payment.**

**1. Coding:** There must be a CPT code or HCPCS code that accurately describes the service performed and/or the drugs provided.


**2. Coverage:** The existence of CPT and/or HCPCS codes used to report the services performed or items furnished does not guarantee coverage.

Medicare only covers a procedure, drug or supply when it is medically necessary. Providers should obtain and follow the policies and guidelines published by Medicare in the Local and National Coverage Determinations.

**3. Payment:** If the proper codes exist and there is coverage established, Medicare must set a payment amount for the drugs, supplies and / or procedures in order for providers to receive payment. Most payment amounts are determined by CMS nationally. There are differences in procedure payment amounts from region to region to reflect geographic differences in provider costs.

**Documentation:** When radiopharmaceuticals or contrast agents are reported, providers must document in the medical record the name of the drug and the amount administered.

Lantheus Medical Imaging cannot guarantee coverage or payment for products or procedures. Payer policies can vary widely. For more specific information, contact the payer directly in order to obtain up to date coverage, coding and payment information.



**2018 Medicare Reimbursement for Nuclear Medicine  
Non-HEU Derived Tc-99m for Medicare Hospital Outpatients<sup>1,2</sup>**

For 2018, CMS will continue the \$10 add-on payment for non-HEU derived Tc-99m for hospital outpatients.

The United States government has established an agenda to eliminate domestic reliance on Tc-99m derived from nuclear reactors using Highly Enriched Uranium (HEU). CMS recognizes that Tc-99m derived from a non-HEU source may have a higher cost. In response, CMS will reimburse providers \$10 per non-HEU derived Tc-99m dose in the hospital outpatient setting in addition to the payment for the imaging procedure.

Under this policy, hospitals report HCPCS code Q9969 (Tc-99m from non-highly enriched uranium source, full cost recovery add-on, per study dose) once per dose along with any diagnostic scan or scans furnished using Tc-99m as long as the Tc-99m doses used can be certified by the hospital to be at least 95 percent derived from non-HEU sources.

1. CMS created HCPCS code Q9969 to report non-HEU Tc-99m doses.

**HCPCS Descriptor**

***Q9969 Tc-99m from non-highly enriched uranium source,  
full cost recovery add-on, per study dose***

2. CMS will reimburse \$10 per dose for Q9969 in addition to the imaging procedure.
3. Hospital reports token \$1 charge per dose for Q9969.

Hospitals do not indicate a dose is from a non-HEU source on their claim form. They simply report HCPCS Q9969 for each non-HEU dose. If asked, a hospital has three options to document a dose was derived from a non-HEU source<sup>2</sup>.

1. Produce invoices, patient dose labels or tracking sheets that indicate that a dose was produced from non-HEU sources.
2. Produce documentation that an entire batch of Tc-99m doses were derived from a non-HEU source for a specified period of time that a single non-HEU generator was in use or manufacturer attestation that a generator is non-HEU generator.
3. If the manufacturer has labeled a generator or a dose attesting to it being derived from a non-HEU source.

If a hospital has any questions about whether they are receiving Tc-99m derived from a non-HEU source, they should contact their radiopharmacy or the generator manufacturer.

For more information, please see Federal Register / Vol. 78, No. 237 / Tuesday, December 10, 2013 p.75002-75003 or Federal Register / Vol. 77, No. 221 / Thursday, November 15, 2012 / p. 68316-68317 or contact your local radiopharmacy or your Tc-99m generator manufacturer.

## 2018 Medicare Reimbursement Information Lantheus Medical Imaging

### Medicare Hospital Outpatient

For 2018, CMS continues to package the payment for diagnostic radiopharmaceuticals, the exercise stress test, CPT 93017, and all pharmacologic stress agents with the SPECT Myocardial Perfusion Imaging (MPI) procedure, CPT 78452, into one single packaged payment.

If a non-HEU derived Tc-99m dose is used, providers will receive a separate add on payment of \$10 per dose by reporting HCPCS code Q9969.

#### Packaged components of HOPPS SPECT Multiple Myocardial Perfusion CPT 78452

Descriptor	2017 <sup>3</sup> payment	2018 <sup>4</sup> payment
78452 SPECT MPI Multiple	\$1138.46	\$1202.60
93017 Exercise test packaged with 78452	\$0 packaged with 78452	\$0 packaged with 78452
Jxxxx Pharmacologic stress agent	\$0 packaged with 78452	\$0 packaged with 78452
A9500 Tc-99m sestamibi	\$0 packaged with 78452	\$0 packaged with 78452
Q9969 non-HEU source Tc-99m per dose	\$10 paid separately	\$10 paid separately

#### Selected 2018 payment Medicare Hospital Outpatients and Physician Office

CPT	Descriptor	APC	Payment HOPPS <sup>4</sup>	Payment Non Hospital Office <sup>5</sup>
78071	Parathyroid planar imaging (including subtraction, when performed); with tomographic (SPECT)	5591	\$349.42	\$377.28
78452	Myocardial Perfusion imaging multiple SPECT	5593	\$1202.60	\$500.39
78582	Pulmonary ventilation (e.g. aerosol or gas) and perfusion imaging	5592	\$453.05	\$352.80
78607	Brain imaging tomographic (SPECT)	5593	\$1202.60	\$369.72
78806	Radiopharmaceutical localization of inflammatory process; whole body	5593	\$1202.60	\$352.44
79101	Radiopharmaceutical therapy, by intravenous administration	5661	\$238.46	\$148.32

Physician Payment amounts as of January 1, 2018. National average payments based on Medicare Conversion factor of \$35.9996 and January, 2018 Medicare Addendum B relative value units. Diagnostic and therapeutic radiopharmaceuticals are reimbursed separately in the non-hospital / office setting based on AWP or invoice. Check with your local Medicare contractor for local payment methodology

**Lantheus Medical Imaging 2018 HCPCS and NDC Information**

**QUADRAMET®** HCPCS code A9604 - NDC 11994-016-01

Samarium 153 lexidronam therapeutic, per treatment dose, up to 150 millicuries.

**Medicare Hospital Outpatients**

- QUADRAMET®- Therapeutic radiopharmaceuticals reimbursed at 106% of ASP.
- Medicare Hospital Outpatient Payment<sup>3</sup> Q4 2017 - A9604, \$13,307.04 per dose.
- Latest payment amount can be seen in Addendum B at:  
<https://www.cms.gov/Medicare/Medicare-Fee-for-Service-Payment/HospitalOutpatientPPS/Addendum-A-and-Addendum-B-Updates.html>.

**Medicare non hospital physician offices / IDTFs**

- QUADRAMET®- Reimbursement based on AWP or invoice, check local Medicare contractor for local payment amount.

Product	HCPCS	NDC codes
Cardiolite® Kit for the Preparation of Technetium Tc-99m Sestamibi for Injection	A9500	<ul style="list-style-type: none"> <li>• NDC 11994-001-20 (20 vials one box)</li> <li>• NDC 11994-001-52 (2 vials one box)</li> <li>• NDC 11994-001-55 (5 vials one box)</li> </ul>
Kit for the Preparation of Technetium Tc-99m Sestamibi for Injection	A9500	<ul style="list-style-type: none"> <li>• NDC 11994-003-20 (20 vials one box)</li> </ul>
Thallous Chloride Thallium 201 Injection	A9505	<ul style="list-style-type: none"> <li>• NDC 11994-427-11</li> <li>• NDC 11994-427-15</li> <li>• NDC 11994-427-19</li> <li>• NDC 11994-427-24</li> <li>• NDC 11994-427-26</li> <li>• NDC 11994-427-28</li> </ul>
Gallium Citrate Ga-67 Injection	A9556	<ul style="list-style-type: none"> <li>• NDC 11994-121-06</li> <li>• NDC 11994-121-08</li> <li>• NDC 11994-121-13</li> <li>• NDC 11994-121-19</li> </ul>
NEUROLITE® Kit for the Preparation of Technetium Tc-99m Bicisate for Injection	A9557	<ul style="list-style-type: none"> <li>• NDC 11994-006-02 (2 kits one package)</li> <li>• NDC 11994-006-05 (5 kits one package)</li> </ul>
Xenon Xe-133 Gas	A9558	<ul style="list-style-type: none"> <li>• NDC 11994-127-11</li> <li>• NDC 11994-127-15</li> <li>• NDC 11994-127-21</li> <li>• NDC 11994-127-25</li> </ul>
Tc-99m from non-highly enriched uranium source, full cost recovery add-on, per study dose	Q9969	<ul style="list-style-type: none"> <li>• Paid \$10 per dose for Tc-99m doses derived from ≥ 95% non-HEU for HOPPS in addition to APC payment for imaging procedure</li> </ul>

## 2018 Medicare Reimbursement Information Lantheus Medical Imaging

### **Citations**

1. Federal Register / Vol. 78, No. 237 / Tuesday, December 10, 2013 p. 75002
2. Federal Register / Vol. 77, No. 221 / Thursday, November 15, 2012 68316
3. <https://www.cms.gov/Medicare/Medicare-Fee-for-Service-Payment/HospitalOutpatientPPS/Hospital-Outpatient-Regulations-and-Notices-Items/CMS-1656-FC.html?DLPage=1&DLEntries=10&DLSort=2&DLSortDir=descending>
4. <https://www.cms.gov/Medicare/Medicare-Fee-for-Service-Payment/HospitalOutpatientPPS/Hospital-Outpatient-Regulations-and-Notices-Items/CMS-1678-FC.html?DLPage=1&DLEntries=10&DLSort=2&DLSortDir=descending>
5. <https://www.cms.gov/Medicare/Medicare-Fee-for-Service-Payment/PhysicianFeeSched/PFS-Federal-Regulation-Notices-Items/CMS-1676-F.html>

NDC codes can be researched at <http://www.accessdata.fda.gov/scripts/cder/ndc/>



331 Treble Cove Road  
N. Billerica, Massachusetts 01862  
USA

# CARDIOLITE®

## Kit for the Preparation of Technetium Tc99m Sestamibi for Injection

FOR DIAGNOSTIC USE

### HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use CARDIOLITE® safely and effectively. See full prescribing information for CARDIOLITE®.

CARDIOLITE®, Kit for the Preparation of Technetium Tc99m Sestamibi for Injection.  
Initial U.S. Approval: December, 1990

#### RECENT MAJOR CHANGES

Use in specific populations (8.3) 10/2007

#### INDICATIONS AND USAGE

CARDIOLITE® is a myocardial perfusion agent indicated for:

- detecting coronary artery disease by localizing myocardial ischemia (reversible defects) and infarction (non-reversible defects)
- evaluating myocardial function and developing information for use in patient management decisions

#### DOSAGE AND ADMINISTRATION

- For Myocardial Imaging: The suggested dose range for I.V. administration of CARDIOLITE® in a single dose to be employed in the average patient (70 Kg) is 370 - 1110 MBq (10 - 30 mCi).
- For Breast Imaging: The recommended dose range for I.V. administration of MIRALUMA® is a single dose of 740 - 1110 MBq (20 - 30 mCi).

#### DOSAGE FORMS AND STRENGTHS

- CARDIOLITE®, Kit for the Preparation of Technetium Tc99m Sestamibi for Injection is supplied as a lyophilized mixture in a 5 mL vial.

#### CONTRAINDICATIONS

- None known

#### WARNINGS AND PRECAUTIONS

- Pharmacologic induction of cardiovascular stress may be associated with serious adverse events such as myocardial infarction, arrhythmia, hypotension, bronchoconstriction and cerebrovascular events.
- CARDIOLITE® has been rarely associated with cutaneous severe allergic and anaphylactic events of angioedema and generalized urticaria. In some patients the allergic symptoms developed on the second injection during CARDIOLITE® imaging.
- Caution should be exercised and emergency equipment should be available when administering CARDIOLITE®.
- Before administering CARDIOLITE® patients should be asked about the possibility of allergic reactions to either drug.
- The contents of the vial are intended only for use in the preparation of Technetium Tc99m Sestamibi and are not to be administered directly to the patient without first undergoing the preparative procedure.

#### ADVERSE REACTIONS

- The following adverse reactions have been reported in ≤ 0.5% of patients: signs and symptoms consistent with seizure occurring shortly after administration of the agent; transient arthritis, angioedema, arrhythmia, dizziness, syncope, abdominal pain, vomiting, and severe hypersensitivity characterized by dyspnea, hypotension, bradycardia, asthenia, and vomiting within two hours after a second injection of Technetium Tc99m Sestamibi. A few cases of flushing, edema, injection site inflammation, dry mouth, fever, pruritis, rash, urticaria and fatigue have also been attributed to administration of the agent.

To report SUSPECTED ADVERSE REACTIONS, contact Lantheus Medical Imaging,

Inc. at 1-800-362-2668 or FDA at 1-800-FDA-1088 or

[www.fda.gov/medwatch](http://www.fda.gov/medwatch).

#### DRUG INTERACTIONS

- Specific drug-drug interactions have not been studied.

#### USE IN SPECIFIC POPULATIONS

- In one study of 46 subjects who received CARDIOLITE® administration, the radioactivity in both children and adolescents exhibited blood PK profiles similar to those previously reported in adults.

#### See 17 FOR PATIENT COUNSELING INFORMATION

Revised: May 2014

513121-0514

### FULL PRESCRIBING INFORMATION: CONTENTS\*

- INDICATIONS AND USAGE
- DOSAGE AND ADMINISTRATION
  - Image Acquisition
  - Radiation Dosimetry
  - Instructions For Preparation
  - Determination of Radiochemical Purity in Technetium Tc99m Sestamibi
- DOSAGE FORMS AND STRENGTHS
- CONTRAINDICATIONS
- WARNINGS AND PRECAUTIONS
  - Warnings
  - General Precautions
- ADVERSE REACTIONS
- DRUG INTERACTIONS
- USE IN SPECIFIC PATIENTS
  - Pregnancy
  - Nursing Mothers

- Pediatric Use
- Geriatric Use
- DRUG ABUSE AND DEPENDENCE
  - Controlled Substance
  - Abuse
  - Dependence
- OVERDOSAGE
- DESCRIPTION
  - Physical Characteristics
  - External Radiation
- CLINICAL PHARMACOLOGY
  - Mechanism of Action
  - Pharmacokinetics
    - Metabolism
    - Elimination
- NONCLINICAL TOXICOLOGY
  - Carcinogenesis, Mutagenesis, Impairment of Fertility
- CLINICAL STUDIES
- REFERENCES
- HOW SUPPLIED/STORAGE AND HANDLING
- PATIENT COUNSELING INFORMATION

\*Sections or subsections omitted from the full prescribing information are not listed.

### 1. INDICATIONS AND USAGE

Myocardial Imaging: CARDIOLITE® Kit for the Preparation of Technetium Tc99m Sestamibi for Injection, is a myocardial perfusion agent that is indicated for detecting coronary artery disease by localizing myocardial ischemia (reversible defects) and infarction (non-reversible defects), in evaluating myocardial function and developing information for use in patient management decisions. CARDIOLITE® evaluation of myocardial ischemia can be accomplished with rest and cardiovascular stress techniques (e.g., exercise or pharmacologic stress in accordance with the pharmacologic stress agent's labeling).

It is usually not possible to determine the age of a myocardial infarction or to differentiate a recent myocardial infarction from ischemia.

Breast Imaging: MIRALUMA®, Kit for the Preparation of Technetium Tc99m Sestamibi for Injection, is indicated for planar imaging as a second line diagnostic drug after mammography to assist in the evaluation of breast lesions in patients with an abnormal mammogram or a palpable breast mass.

MIRALUMA® is not indicated for breast cancer screening, to confirm the presence or absence of malignancy, and it is not an alternative to biopsy.

### 2. DOSAGE AND ADMINISTRATION

For Myocardial Imaging: The suggested dose range for I.V. administration of CARDIOLITE® in a single dose to be employed in the average patient (70 Kg) is 370 - 1110 MBq (10 - 30 mCi).

For Breast Imaging: The recommended dose range for I.V. administration of MIRALUMA® is a single dose of 740 - 1110 MBq (20 - 30 mCi).

#### 2.1 Image Acquisition

Breast Imaging: It is recommended that images are obtained with a table overlay to separate breast tissue from the myocardium and liver, and to exclude potential activity that may be present in the opposite breast. For lateral images, position the patient prone with the ipsilateral arm comfortably above the head, shoulders flat against the table, head turned to the side and relaxed, with the breast imaged pendent through an overlay cutout. The breast should not be compressed on the overlay. For anterior images, position the patient supine with both arms behind the head. For either lateral or anterior images, shield the chest and abdominal organs, or remove them from the field of view.

For complete study, sets of images should be obtained five minutes after the injection, and in the following sequence:

Beginning five minutes after the injection of Technetium Tc99m Sestamibi:

- ten-minute lateral image of breast with abnormality
- ten-minute lateral image of contralateral breast
- ten-minute anterior image of both breasts

#### 2.2 Radiation Dosimetry

The radiation doses to organs and tissues of an average patient (70 Kg) per 1110 MBq (30 mCi) of Technetium Tc99m Sestamibi injected intravenously are shown in Table 1.0.

Table 1.0. Radiation Absorbed Doses from Tc99m Sestamibi

Organ	Estimated Radiation Absorbed Dose			
	REST			
	2.0 hour void		4.8 hour void	
rads/ 30 mCi	mGy/ 1110 MBq	rads/ 30 mCi	mGy/ 1110 MBq	
Breasts	0.2	2.0	0.2	1.9
Gallbladder Wall	2.0	20.0	2.0	20.0
Small Intestine	3.0	30.0	3.0	30.0
Upper Large Intestine Wall	5.4	55.5	5.4	55.5
Lower Large Intestine Wall	3.9	40.0	4.2	41.1
Stomach Wall	0.6	6.1	0.6	5.8
Heart Wall	0.5	5.1	0.5	4.9
Kidneys	2.0	20.0	2.0	20.0
Liver	0.6	5.8	0.6	5.7
Lungs	0.3	2.8	0.3	2.7
Bone Surfaces	0.7	6.8	0.7	6.4
Thyroid	0.7	7.0	0.7	7.0
Ovaries	1.5	15.5	1.6	15.5
Testes	0.3	3.4	0.4	3.9
Red Marrow	0.5	5.1	0.5	5.0
Urinary Bladder Wall	2.0	20.0	4.2	41.1
Total Body	0.5	4.8	0.5	4.8

### STRESS

Organ	STRESS			
	2.0 hour void		4.8 hour void	
	rads/ 30 mCi	mGy/ 1110 MBq	rads/ 30 mCi	mGy/ 1110 MBq
Breasts	0.2	2.0	0.2	1.8
Gallbladder Wall	2.8	28.9	2.8	27.8
Small Intestine	2.4	24.4	2.4	24.4
Upper Large Intestine Wall	4.5	44.4	4.5	44.4
Lower Large Intestine Wall	3.3	32.2	3.3	32.2
Stomach Wall	0.6	5.3	0.5	5.2
Heart Wall	0.5	5.6	0.5	5.3
Kidneys	1.7	16.7	1.7	16.7
Liver	0.4	4.2	0.4	4.1
Lungs	0.3	2.6	0.2	2.4
Bone Surfaces	0.6	6.2	0.6	6.0
Thyroid	0.3	2.7	0.2	2.4
Ovaries	1.2	12.2	1.3	13.3
Testes	0.3	3.1	0.3	3.4
Red Marrow	0.5	4.6	0.5	4.4
Urinary Bladder Wall	1.5	15.5	3.0	30.0
Total Body	0.4	4.2	0.4	4.2

Radiation dosimetry calculations performed by Radiation Internal Dose Information Center, Oak Ridge Institute for Science and Education, PO Box 117, Oak Ridge, TN 37831-0117.

### 2.3 Instructions For Preparation

Preparation of the Technetium Tc99m Sestamibi from the Kit for the Preparation of Technetium Tc99m Sestamibi is done by the following aseptic procedure:

#### General Procedure:

- Prior to adding the Sodium Pertechnetate Tc99m Injection to the vial, inspect the vial carefully for the presence of damage, particularly cracks, and do not use the vial if found. Tear off a radiation symbol and attach it to the neck of the vial.
- Waterproof gloves should be worn during the preparation procedure. Remove the plastic disc from the vial and swab the top of the vial closure with alcohol to sanitize the surface.

#### Boiling Water Bath Procedure:

- Place the vial in a suitable radiation shield with a fitted radiation cap.
- With a sterile shielded syringe, aseptically obtain additive-free, sterile, non-pyrogenic Sodium Pertechnetate Tc99m Injection [925 - 5550 MBq, (25 - 150 mCi)] in approximately 1 to 3 mL.
- Aseptically add the Sodium Pertechnetate Tc99m Injection to the vial in the lead shield. Without withdrawing the needle, remove an equal volume of headspace to maintain atmospheric pressure within the vial.
- Shake vigorously, about 5 to 10 quick upward-downward motions.
- Remove the vial from the lead shield and place upright in an appropriately shielded and contained boiling water bath, such that the vial is suspended above the bottom of the bath, and boil for 10 minutes. Timing for 10 minutes is begun as soon as the water begins to boil again. Do not allow the boiling water to come in contact with the aluminum crimp.
- Remove the vial from the water bath, place in the lead shield and allow to cool for fifteen minutes.

#### Recon-o-Stat (thermal cycler) Procedure:

- Place the vial in the thermal cycler radiation shield.
- With a sterile shielded syringe, aseptically obtain additive-free, sterile, non-pyrogenic Sodium Pertechnetate Tc99m Injection [925 - 5550 MBq, (25 - 150 mCi)] in approximately 1 to 3 mL.
- Aseptically add the Sodium Pertechnetate Tc99m Injection to the vial in the lead shield. Without withdrawing the needle, remove an equal volume of headspace to maintain atmospheric pressure within the vial.
- Shake vigorously, about 5 to 10 quick upward-downward motions.
- Place shield on sample block. While slightly pressing downward, give the shield a quarter turn to make certain there is a firm fit between the shield and the sample block.
- Press the proceed button to initiate the program (the thermal cycler automatically heats & cools the vial and contents). Please see the Recon-o-Stat Instruction Manual for further details.

#### General Procedure (cont.):

- Using proper shielding, the vial contents should be visually inspected. Use only if the solution is clear and free of particulate matter and discoloration.
- Assay the reaction vial using a suitable radioactivity calibration system. Record the Technetium Tc99m concentration, total volume, assay time and date, expiration time and lot number on the vial shield label and affix the label to the shield.
- Store the reaction vial containing the Technetium Tc99m Sestamibi at 15° to 25°C (59-77°F) until use; at such time the product should be aseptically withdrawn. Technetium Tc99m Sestamibi should be used within six hours of preparation. The vial contains no preservative.

**Note:** Adherence to the above product reconstitution instructions is recommended.

The potential for cracking and significant contamination exists whenever vials containing radioactive material are heated.

Product should be used within 6 hours after preparation. Final product with radiochemical purity of at least 90% was used in the clinical trials that established safety and effectiveness. The radiochemical purity was determined by the following method.




## 2.4 Determination of Radiochemical Purity in Technetium Tc99m Sestamibi

- Obtain a Baker-Flex Aluminum Oxide coated, plastic TLC plate, #1 B-F, pre-cut to 2.5 cm x 7.5 cm.
- Dry the plate or plates at 100°C for 1 hour and store in a desiccator. Remove pre-dried plate from the desiccator just prior to use.
- Apply 1 drop of ethanol\* using a 1 mL syringe with a 22-26 gauge needle, 1.5 cm from the bottom of the plate. THE SPOT SHOULD NOT BE ALLOWED TO DRY.
- Add 2 drops of Technetium Tc99m Sestamibi solution, side by side on top of the ethanol\* spot. Return the plate to a desiccator and allow the sample spot to dry (typically 15 minutes).
- The TLC tank is prepared by pouring ethanol\* to a depth of 3-4 mm. Cover the tank and let it equilibrate for ~10 minutes.
- Develop the plate in the covered TLC tank in ethanol\* for a distance of 5 cm from the point of application.
- Cut the TLC plate 4 cm from the bottom and measure the Tc99m activity in each piece by appropriate radiation detector.
- Calculate the % Tc99m Sestamibi as:

$$\% \text{ Tc99m Sestamibi} = \frac{\mu\text{Ci Top Piece}}{\mu\text{Ci Both Pieces}} \times 100$$

Figure 1.0 TLC Plate Diagram



\*The ethanol used in this procedure should be 95% or greater. Absolute ethanol (99%) should remain at  $\geq 95\%$  ethanol content for one week after opening if stored tightly capped, in a cool dry place.

## 3. DOSAGE FORMS AND STRENGTHS

CARDIOLITE<sup>®</sup>, Kit for the Preparation of Technetium Tc99m Sestamibi for Injection is supplied as a lyophilized mixture in a 5 mL vial.

## 4. CONTRAINDICATIONS

None known.

## 5. WARNINGS AND PRECAUTIONS

### 5.1 Warnings

In studying patients in whom cardiac disease is known or suspected, care should be taken to assure continuous monitoring and treatment in accordance with safe, accepted clinical procedure. Infrequently, death has occurred 4 to 24 hours after Tc99m Sestamibi use and is usually associated with exercise stress testing (See Section 5.2).

Pharmacologic induction of cardiovascular stress may be associated with serious adverse events such as myocardial infarction, arrhythmia, hypotension, bronchoconstriction and cerebrovascular events. Caution should be used when pharmacologic stress is selected as an alternative to exercise; it should be used when indicated and in accordance with the pharmacologic stress agent's labeling.

Technetium Tc99m Sestamibi has been rarely associated with acute severe allergic and anaphylactic events of angioedema and generalized urticaria. In some patients the allergic symptoms developed on the second injection during CARDIOLITE<sup>®</sup> imaging. Patients who receive CARDIOLITE<sup>®</sup> or MIRALUMA<sup>®</sup> imaging are receiving the same drug. Caution should be exercised and emergency equipment should be available when administering Technetium Tc99m Sestamibi. Also, before administering either CARDIOLITE<sup>®</sup> or MIRALUMA<sup>®</sup>, patients should be asked about the possibility of allergic reactions to either drug.

### 5.2 General Precautions

The contents of the vial are intended only for use in the preparation of Technetium Tc99m Sestamibi and are not to be administered directly to the patient without first undergoing the preparative procedure.

Radioactive drugs must be handled with care and appropriate safety measures should be used to minimize radiation exposure to clinical personnel. Also, care should be taken to minimize radiation exposure to the patients consistent with proper patient management.

Contents of the kit before preparation are not radioactive. However, after the Sodium Pertechnetate Tc99m Injection is added, adequate shielding of the final preparation must be maintained. The components of the kit are sterile and non-pyrogenic. It is essential to follow directions carefully and to adhere to strict aseptic procedures during preparation.

Technetium Tc99m labeling reactions depend on maintaining the stannous ion in the reduced state. Hence, Sodium Pertechnetate Tc99m Injection containing oxidants should not be used.

Technetium Tc99m Sestamibi should not be used more than six hours after preparation.

Radiopharmaceuticals should be used only by physicians who are qualified by training and experience in the safe use and handling of radionuclides and whose experience and training have been approved by the appropriate government agency authorized to license the use of radionuclides.

Stress testing should be performed only under the supervision of a qualified physician and in a laboratory equipped with appropriate resuscitation and support apparatus.

The most frequent exercise stress test endpoints sufficient to stop the test reported during controlled studies (two-thirds were cardiac patients) were:

Fatigue	35%
Dyspnea	17%
Chest Pain	16%
ST-depression	7%
Arrhythmia	1%

## 6. ADVERSE REACTIONS

Adverse events were evaluated in 3741 adults who were evaluated in clinical studies. Of these patients, 3068 (77% men, 22% women, and 0.7% of the patient's genders were not recorded) were in cardiac clinical trials and 673 (100% women) in breast imaging trials. Cases of angina, chest pain, and death have occurred (see Section 5). Adverse events reported at a rate of 0.5% or greater after receiving Technetium Tc99m Sestamibi administration are shown in the following table:

Table 2.0 Selected Adverse Events Reported in > 0.5% of Patients Who Received Technetium Tc99m Sestamibi in Either Breast or Cardiac Clinical Studies*				
Body System	Breast Studies		Cardiac Studies	
	Women n = 673	Women n = 685	Men n = 2361	Total n = 3046
Body as a Whole	21 (3.1%)	6 (0.9%)	17 (0.7%)	23 (0.8%)
Headache	11 (1.6%)	2 (0.3%)	4 (0.2%)	6 (0.2%)
Cardiovascular	9 (1.3%)	24 (3.5%)	75 (3.2%)	99 (3.3%)
Chest Pain/Angina	0 (0%)	18 (2.6%)	46 (1.9%)	64 (2.1%)
ST segment changes	0 (0%)	11 (1.6%)	29 (1.2%)	40 (1.3%)
Digestive System	8 (1.2%)	4 (0.6%)	9 (0.4%)	13 (0.4%)
Nausea	4 (0.6%)	1 (0.1%)	2 (0.1%)	3 (0.1%)
Special Senses	132 (19.6%)	62 (9.1%)	160 (6.8%)	222 (7.3%)
Taste Perversion	129 (19.2%)	60 (8.8%)	157 (6.6%)	217 (7.1%)
Parosmia	8 (1.2%)	6 (0.9%)	10 (0.4%)	16 (0.5%)

\*Excludes the 22 patients whose gender was not recorded.

In the clinical studies for breast imaging, breast pain was reported in 12 (1.7%) of the patients. In 11 of these patients the pain appears to be associated with biopsy/surgical procedures.

The following adverse reactions have been reported in  $\leq 0.5\%$  of patients: signs and symptoms consistent with seizure occurring shortly after administration of the agent; transient arthritis, angioedema, arrhythmia, dizziness, syncope, abdominal pain, vomiting, and severe hypersensitivity characterized by dyspnea, hypotension, bradycardia, asthenia, and vomiting within two hours after a second injection of Technetium Tc99m Sestamibi. A few cases of flushing, edema, injection site inflammation, dry mouth, fever, pruritis, rash, urticaria and fatigue have also been attributed to administration of the agent.

## 7. DRUG INTERACTIONS

Specific drug-drug interactions have not been studied.

## 8. USE IN SPECIFIC PATIENTS

### 8.1 Pregnancy

Animal reproduction and teratogenicity studies have not been conducted with Technetium Tc99m Sestamibi. It is also not known whether Technetium Tc99m Sestamibi can cause fetal harm when administered to a pregnant woman or can affect reproductive capacity. There have been no studies in pregnant women. Technetium Tc99m Sestamibi should be given to a pregnant woman only if clearly needed.

### 8.3 Nursing Mothers

Technetium Tc99m Pertechnetate is excreted in human milk during lactation. It is not known whether Technetium Tc99m Sestamibi is excreted in human milk. Therefore, formula feedings should be substituted for breast feedings.

### 8.4 Pediatric Use

Safety and effectiveness in the pediatric population have not been established.

No evidence of diagnostic efficacy or clinical utility of CARDIOLITE<sup>®</sup> scan was found in clinical studies of children and adolescents with Kawasaki disease.

A prospective study of 445 pediatric patients with Kawasaki disease was designed to determine the predictive value of CARDIOLITE<sup>®</sup> rest and stress myocardial perfusion imaging to define a pediatric population with Kawasaki disease that was at risk of developing cardiac events. Cardiac events were defined as cardiac death, MI, hospitalization due to cardiac etiology, heart failure, CABG or coronary angioplasty. The standard of truth was defined as cardiac events occurring 6 months following the administration of CARDIOLITE<sup>®</sup>. Only three cardiac events were observed at six months in this study. In all three cases, the scan was negative. No clinically meaningful measurements of sensitivity, specificity or other diagnostic performance parameters could be demonstrated in this study.

A ten year retrospective case history study of pediatric Kawasaki disease patients who completed CARDIOLITE<sup>®</sup> myocardial perfusion imaging and who had coronary angiography within three months of the CARDIOLITE<sup>®</sup> scan was designed to measure sensitivity and specificity of CARDIOLITE<sup>®</sup> scan. Out of 72 patients who had both evaluable CARDIOLITE<sup>®</sup> scans and evaluable angiographic images, only one patient had both an abnormal angiogram and an abnormal CARDIOLITE<sup>®</sup> scan. No clinically meaningful measurements of sensitivity, specificity or other diagnostic performance parameters could be demonstrated in this study.

In a clinical pharmacology study, 46 pediatric patients with Kawasaki disease received CARDIOLITE<sup>®</sup> administration at the following doses: 0.1 - 0.2 mCi/kg for rest, 0.3 mCi/kg for stress in one day studies; 0.2 mCi/kg for rest and 0.2 mCi/kg for stress in two day studies.

The radioactivity both in younger children and in adolescents exhibited PK profiles similar to those previously reported in adults (See Section 12).

The radiation absorbed doses in adolescents, both at rest and stress, were similar to those observed in adults (see Section 2). When comparing weight-adjusted radioactivity (up to 0.3 mCi/kg) doses administered to adolescents and younger children to the recommended dose administered to adults (up to 30 mCi), the radiation absorbed doses in both adolescents and younger children were similar to those in adults.

Adverse events were evaluated in 609 pediatric patients from the three clinical studies described above. The frequency and the type of the adverse events were similar to the ones observed in the studies of CARDIOLITE<sup>®</sup> in adults. Two of the 609 had a serious adverse event: one patient received a CARDIOLITE<sup>®</sup> overdose but remained asymptomatic, and one patient had an asthma exacerbation following administration.

## 8.5 Geriatric Use

Of 3068 patients in clinical studies of CARDIOLITE<sup>®</sup>, Kit for the Preparation of Technetium Tc99m Sestamibi for Injection, 693 patients were 65 or older and 121 were 75 or older.

Of 673 patients in clinical studies of MIRALUMA<sup>®</sup>, Kit for the Preparation of Technetium Tc99m Sestamibi for Injection, 138 patients were 65 or older and 30 were 75 or older.

Based on the evaluation of the frequency of adverse events and review of vital signs data, no overall differences in safety were observed between these subjects and younger subjects. Although reported clinical experience has not identified differences in response between elderly and younger patients, greater sensitivity of some older individuals cannot be ruled out.

## 9. DRUG ABUSE AND DEPENDENCE

### 9.1 Controlled Substance

Not applicable.

### 9.2 Abuse

Not applicable.

### 9.3 Dependence

Not applicable.

## 10. OVERDOSAGE

The clinical consequences of overdosing with CARDIOLITE<sup>®</sup> are not known.

## 11. DESCRIPTION

Each 5 mL vial contains a sterile, non-pyrogenic, lyophilized mixture of:

- Tetrakis (2-methoxy isobutyl isonitrile) Copper (I) tetrafluoroborate - 1.0 mg
- Sodium Citrate Dihydrate - 2.6 mg
- L-Cysteine Hydrochloride Monohydrate - 1.0 mg
- Mannitol - 20 mg
- Stannous Chloride, Dihydrate, minimum (SnCl<sub>2</sub>•2H<sub>2</sub>O) - 0.025 mg
- Stannous Chloride, Dihydrate, (SnCl<sub>2</sub>•2H<sub>2</sub>O) - 0.075 mg
- Tin Chloride (stannous and stannic) Dihydrate, maximum (as SnCl<sub>2</sub>•2H<sub>2</sub>O) - 0.086 mg

Prior to lyophilization the pH is 5.3 to 5.9. The contents of the vial are lyophilized and stored under nitrogen.

This drug is administered by intravenous injection for diagnostic use after reconstitution with sterile, non-pyrogenic, oxidant-free Sodium Pertechnetate Tc99m Injection. The pH of the reconstituted product is 5.5 (5.0 - 6.0). No bacteriostatic preservative is present.

The precise structure of the technetium complex is Tc99m[MIBI]<sub>6</sub><sup>+</sup> where MIBI is 2-methoxy isobutyl isonitrile.

### 11.1 Physical Characteristics

Technetium Tc99m decays by isomeric transition with a physical half-life of 6.02 hours<sup>1</sup>. Photons that are useful for detection and imaging studies are listed below in Table 3.0.

Table 3.0. Principal Radiation Emission Data

Radiation	Mean %/ Disintegration	Mean Energy (KeV)
Gamma -2	89.07	140.5

<sup>1</sup>Kocher, David, C., Radioactive Decay Data Tables, DOE/TIC-11026, 108(1981).

### 11.2 External Radiation

The specific gamma ray constant for Tc99m is 5.4 microcoulombs/Kg-MBq-hr (0.78R/mCi-hr) at 1 cm. The first half value layer is 0.017 cm of Pb. A range of values for the relative attenuation of the radiation emitted by this radionuclide that results from interposition of various thicknesses of Pb is shown in Table 4.0. To facilitate control of the radiation exposure from Megabequerel (millicurie) amounts of this radionuclide, the use of a 0.25 cm thickness of Pb will attenuate the radiation emitted by a factor of 1,000.

Table 4.0. Radiation Attenuation by Lead Shielding

Shield Thickness (Pb) cm	Coefficient of Attenuation
0.017	0.5
0.08	10 <sup>-1</sup>
0.16	10 <sup>-2</sup>
0.25	10 <sup>-3</sup>
0.33	10 <sup>-4</sup>

To correct for physical decay of this radionuclide, the fractions that remain at selected intervals after the time of calibration are shown in Table 5.0.

**Table 5.0. Physical Decay Chart; Tc99m Half-Life 6.02 Hours**

Hours	Fraction Remaining	Hours	Fraction Remaining
0*	1.000	8	.398
1	.891	9	.355
2	.794	10	.316
3	.708	11	.282
4	.631	12	.251
5	.562		
6	.501		
7	.447		

\*Calibration Time

**12. CLINICAL PHARMACOLOGY**

**12.1 Mechanism of Action**

Technetium Tc99m Sestamibi is a cationic Tc99m complex which has been found to accumulate in viable myocardial tissue in a manner analogous to that of thallous chloride TI-201. Scintigraphic images obtained in humans after the intravenous administration of the drug have been comparable to those obtained with thallous chloride TI-201 in normal and abnormal myocardial tissue.

Animal studies have shown that myocardial uptake is not blocked when the sodium pump mechanism is inhibited. Although studies of subcellular fractionation and electron micrographic analysis of heart cell aggregates suggest that Tc99m Sestamibi cellular retention occurs specifically within the mitochondria as a result of electrostatic interactions, the clinical relevance of these findings has not been determined.

The mechanism of Tc99m Sestamibi localization in various types of breast tissue (e.g., benign, inflammatory, malignant, fibrous) has not been established.

**12.3 Pharmacokinetics**

Pulmonary activity is negligible even immediately after injection. Blood clearance studies indicate that the fast clearing component clears with a  $t_{1/2}$  of 4.3 minutes at rest, and clears with a  $t_{1/2}$  of 1.6 minutes under exercise conditions. At five minutes post injection about 8% of the injected dose remains in circulation. There is less than 1% protein binding of Technetium Tc99m Sestamibi in plasma. The myocardial biological half-life is approximately six hours after a rest or exercise injection. The biological half-life for the liver is approximately 30 minutes after a rest or exercise injection. The effective half-life of clearance (which includes both the biological half-life and radionuclide decay) for the heart is approximately 3 hours, and for the liver is approximately 30 minutes, after a rest or exercise injection. The ideal imaging time reflects the best compromise between heart count rate and surrounding organ uptake.

Myocardial uptake which is coronary flow dependent is 1.2% of the injected dose at rest and 1.5% of the injected dose at exercise. Table 6.0 illustrates the biological clearance as well as effective clearance (which includes biological clearance and radionuclide decay) of Tc99m Sestamibi from the heart and liver.

[Organ concentrations expressed as percentage of injected dose; data based on an average of 5 subjects at rest and 5 subjects during exercise].

**Table 6.0 Biological and Effective Clearance**

Time	REST				STRESS			
	Heart		Liver		Heart		Liver	
	Biological	Effective	Biological	Effective	Biological	Effective	Biological	Effective
5 min.	1.2	1.2	19.6	19.4	1.5	1.5	5.9	5.8
30 min.	1.1	1.0	12.2	11.5	1.4	1.3	4.5	4.2
1 hour	1.0	0.9	5.6	5.0	1.4	1.2	2.4	2.1
2 hours	1.0	0.8	2.2	1.7	1.2	1.0	0.9	0.7
4 hours	0.8	0.5	0.7	0.4	1.0	0.6	0.3	0.2

A study in a dog myocardial ischemia model reported that Technetium Tc99m Sestamibi undergoes myocardial distribution (redistribution), although more slowly and less completely than thallous chloride TI-201. A study in a dog myocardial infarction model reported that the drug showed no redistribution of any consequence. Definitive human studies to demonstrate possible redistribution have not been reported. In patients with documented myocardial infarction, imaging revealed the infarct up to four hours post dose.

**12.3.1 Metabolism**

The agent is excreted without any evidence of metabolism.

**12.3.2 Elimination**

The major pathway for clearance of Tc99m Sestamibi is the hepatobiliary system. Activity from the gall bladder appears in the intestines within one hour of injection. Twenty-seven percent of the injected dose is excreted in the urine, and approximately thirty-three percent of the injected dose is cleared through the feces in 48 hours.

**13. NONCLINICAL TOXICOLOGY**

**13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility**

In comparison with most other diagnostic technetium labeled radiopharmaceuticals, the radiation dose to the ovaries (1.5 rads/30 mCi at rest, 1.2 rads/30 mCi at exercise) is high. Minimal exposure (ALARA) is necessary in women of childbearing capability. (See Section 2.)

The active intermediate, Cu(MIBI)<sub>2</sub>BF<sub>6</sub>, was evaluated for genotoxic potential in a battery of five tests. No genotoxic activity was observed in the Ames, CHO/HPRT and sister chromatid exchange tests (all in vitro). At cytotoxic concentrations (> 20 µg/mL), an increase in cells with chromosome aberrations was observed in the in vitro human lymphocyte assay. Cu(MIBI)<sub>2</sub>BF<sub>6</sub> did not show genotoxic effects in the in vivo mouse micronucleus test at a dose which caused systemic and bone marrow toxicity (9 mg/kg, > 600 X maximal human dose).

**14. CLINICAL STUDIES**

**CLINICAL TRIALS:**

**MYOCARDIAL IMAGING:** In a trial of rest and stress CARDIOLITE® imaging, the relationship of normal or abnormal perfusion scans and long term cardiac events was evaluated in 521 patients (511 men, 10 women) with stable chest pain. There were

73.9% Caucasians, 25.9% Blacks and 0.2% Asians. The mean age was 59.6 years (range: 29 to 84 years). All patients had a baseline rest and exercise CARDIOLITE® scan and were followed for 13.2 ± 4.9 months (range: 1 to 24 months). Images were correlated with the occurrence of a cardiac event (cardiac death or non-fatal myocardial infarction). In this trial as summarized in Table 7.0, 24/521 (4.6%) had a cardiac event.

**Table 7.0**

Cardiac Events			
Baseline Scan <sup>(a)</sup>	Proportion of patients with events by scan results <sup>(a)</sup>	Proportion of scan result in patients with events; N=24 <sup>(a)</sup>	Proportion of event-free patients by scan result <sup>(a)</sup>
Normal	1/206 (0.5%)	1/24 (4.2%)	205/206 (99.5%)
Abnormal	23/315 (7.3%) <sup>(b)</sup>	23/24 (95.8%) <sup>(b)</sup>	292/315 (92.7%) <sup>(b)</sup>

(a) Note: Similar findings were found in two studies with patients who had pharmacologic stress CARDIOLITE® imaging.

(b) p<0.01

Although patients with normal images had a lower cardiac event rate than those with abnormal images, in all patients with abnormal images it was not possible to predict which patient would be likely to have further cardiac events; i.e., such individuals were not distinguishable from other patients with abnormal images.

The findings were not evaluated for defect location, disease duration, specific vessel involvement or intervening management.

In earlier trials, using a template consisting of the anterior wall, inferior-posterior wall and isolated apex, localization in the anterior or inferior-posterior wall in patients with suspected angina or coronary artery disease was shown. Disease localization isolated to the apex has not been established. In adults, Tc99m Sestamibi has not been studied or evaluated in cardiac disorders other than coronary artery disease.

**BREAST IMAGING:** MIRALUMA® was evaluated in two multicenter, clinical trials of a total of 673 woman patients. Overall the mean age was 52 (range 23 to 87 years). The racial and ethnic representation was 70% Caucasian, 15% African-American, 14% Hispanic and 1% Asian.

Both clinical studies evaluated women who were referred for further evaluation for either: 1) a mammographically detected (with varying degrees of malignant likelihood) but not palpable breast lesion (study A, n=387, mean age = 54 years), or 2) a palpable breast lesion (study B, n=286, mean age = 50 years). In both studies all patients were scheduled for biopsy.

MIRALUMA® (20 - 30 mCi) was injected intravenously in a vein that was contralateral to the breast lesion in question. Planar imaging was completed with a high resolution collimator with a 10% window centered at 140 KeV, and 128 x 128 matrix. An initial marker image, that was not used in the data analysis, was obtained using a cobalt Co57 point source as a marker of a palpable mass. Images were obtained 5 minutes after injection as follows: lateral image of the affected breast for 10 minutes, lateral image of the contralateral breast for 10 minutes, and an anterior image of both breasts for 10 minutes. For the lateral image the patients were positioned in a prone position. For the anterior image, the patients were supine. The MIRALUMA® scintigraphic images were read in a randomized method by two groups of three blinded readers. MIRALUMA® uptake was scored as: normal (no uptake), equivocal, low, moderate, or high uptake. The results of MIRALUMA® images and mammography were analyzed in comparison to histopathologic findings of malignant or non-malignant disease.

As shown in Table 8.0 for the 483 evaluable patients, the sensitivity and specificity of any degree of MIRALUMA® uptake appear to vary with the presence or absence of palpable mass.

**TABLE 8.0 Overall MIRALUMA® Blinded Results of Target Lesions<sup>(a)</sup> Identified at Study Entry<sup>(b)</sup>**

STATISTIC	Study A Non-Palpable Mass and an Abnormal Mammogram	Study B Palpable Mass
Number of Patients and Lesions	N=277 Patients with 300 Lesions	N=206 Patients with 240 Lesions
Sensitivity	52(42,62) <sup>(c)</sup>	76(67,83)
Specificity	94(89,96)	85(77,91)
PPV <sup>(d)</sup>	79(67,88)	83(74,89)
NPV <sup>(d)</sup>	80(74,85)	78(69,84)
Agreement	80(75,85)	80(75,85)
Prevalence	32(27,37)	49(43,56)
(a)	Excludes all discordant lesions not identified at entry and excludes 25 equivocal interpretations from Study A and 32 equivocal interpretations from Study B (see Tables 9.0 and 10.0)	
(b)	Some patients had more than one target lesion	
(c)	Median and approximated 95% Confidence Interval	
(d)	PPV= Positive Predict Value; NPV= Negative Predict Value	

In a separate retrospective subset analyses of 259 patients with dense (heterogeneously/extremely dense) and 275 patients with fatty (almost entirely fat/numerous vague densities) breast tissue, the MIRALUMA® results were similar. Overall, the studies were not designed to compare the performance of MIRALUMA® with the performance of mammography in patients with breast densities or other coexistent breast tissue disorders.

In general the histology seems to correlate with the degree of MIRALUMA® uptake. As shown in Tables 9.0 and 10.0, the majority of the normal MIRALUMA® images are associated with non-malignant tissue (78-81%) and the majority of low, moderate or high uptake MIRALUMA® images are associated with malignant disease (79-83%). In an individual patient, however, the intensity of MIRALUMA® uptake can not be used to confirm the presence or absence of malignancy. Equivocal results do not have a correlation with histology.

**TABLE 9.0 Degree of MIRALUMA® Breast Imaging Uptake in Comparison to Histopathology Results in Patients with Mammographically Detected Non-Palpable Lesions\* (Study A)**

	Normal Uptake N = 249 lesions	Equivocal Uptake N = 25 lesions	Low, Moderate or High Uptake N = 66 lesions
Non-malignant**	201 (81%)	14 (56%)	14 (21%)
Malignant	48 (19%)	11 (44%)	52 (79%)

\* Median finding for 3 blinded readers

\*\* Includes benign tissue, fibroadenoma, benign intramammary nodes, radial scar.

**TABLE 10.0 Degree of MIRALUMA® Breast Imaging Uptake in Comparison to Histopathology Results in Patients with Palpable Lesions\* (Study B)**

	Normal Uptake N = 129 lesions	Equivocal Uptake N = 32 lesions	Low, Moderate or High Uptake N = 115 lesions
Non-malignant**	100 (78%)	19 (59%)	20 (17%)
Malignant	29 (22%)	13 (41%)	95 (83%)

\* Median finding for 3 blinded readers

\*\* Includes benign tissue, fibroadenoma, benign intramammary nodes, radial scar.

An estimate of the likelihood of malignancy based on the MIRALUMA® uptake score in combination with the mammographic score has not been studied.

In these two studies approximately 150 additional, non-biopsied lesions were found to be positive after MIRALUMA® imaging. These lesions were identified in sites that did not physically correlate with identified entry criteria mammographic lesions and these lesions were not palpable. These lesions were not biopsied. Whether these lesions were benign or malignant is not known. MIRALUMA® uptake can occur in both benign and malignant disease. THE CLINICAL USEFULNESS OF A POSITIVE MIRALUMA® IMAGE IN THE ABSENCE OF AN ABNORMAL MAMMOGRAM OR A PALPABLE LESION IS NOT KNOWN.

**15. REFERENCES**

Not applicable.

**16. HOW SUPPLIED/STORAGE AND HANDLING**

CARDIOLITE®, Kit for the Preparation of Technetium Tc99m Sestamibi for Injection is supplied as a 5 mL vial in kits of five (5) vials (NDC # 11994-001-55) and twenty (20) vials (NDC # 11994-001-20), sterile and non-pyrogenic.

The patient dose should be measured by a suitable radioactivity calibration system immediately prior to patient administration. Radiochemical purity should be checked prior to patient administration.

Parenteral drug products should be inspected visually for particulate matter and discoloration prior to administration whenever solution and container permit.

Prior to lyophilization the pH is between 5.3-5.9. The contents of the vial are lyophilized and stored under nitrogen. Store at 15-25°C (59-77° F) before and after reconstitution.

Technetium Tc99m Sestamibi contains no preservatives. Included in each five (5) vial kit is one (1) package insert, six (6) vial shield labels and six (6) radiation warning labels. Included in each twenty (20) vial kit is one (1) package insert, twenty four (24) vial shield labels and twenty four (24) radiation warning labels.

This reagent kit is approved for distribution to persons licensed pursuant to the Code of Massachusetts Regulations 105 CMR 120.500 for the uses listed in 105 CMR 120.547 or 120.552, or under equivalent regulations of the U.S. Nuclear Regulatory Commission, Agreement States or Licensing States.

**17. PATIENT COUNSELING INFORMATION**

CARDIOLITE® and MIRALUMA® are different names for the same drug. Patients should be advised to inform their health care provider if they had an allergic reaction to either drug or if they had an imaging study with either drug.



(For Massachusetts and International call 978-667-9531)



[www.lantheus.com](http://www.lantheus.com)

Cardiolite, TechneLite, NEUROLITE and corporate logo display are registered trademarks of Lantheus Medical Imaging, Inc.  
 ©2017 Lantheus Medical Imaging, Inc. All rights reserved. LNMR-0117